

## PERFORMANCE DATA SHEET

2135

### Human CD274(B7-H1)-muIg Fusion Protein\*

For maximal recovery of contents  
please quick spin vial before opening

**CATALOG#: 541-820 (Preservative-free)**

**QUANTITY: 25 µg**

**CONCENTRATION: 0.5 mg/ml**

**Molecular Structure:** A soluble fusion protein consisting of the murine CD8 alpha leader sequence, the mature extracellular (224aa) domain of human CD274 fused to murine IgG2a Fc + hinge (233aa).

**muCD8 apha signal peptide residual amino acids+ linker:** (1) kqapelrgsas

**CD274 mature EC: (224aa):**

(13)ftvtvpkdlyvveygsnmtieckfpvekqldlaalivywemedkniiqfvhgeedlkvqhssyrqrarllkdqslgnaalqitdvklqdagvyrcmisyyggadykritvkvnapynkinqrilv  
vdpvtseheltcqaegy pkaevitwssdhqvlsgkttttnskreeklfnvtstlrntttneifyctfrldpeenhtaelvipelplahppnerthtr

**Linker +Murine IgG2a Hinge + Fc (235 aa):**

(237)gteprgptikpcppckcpapnllggpsvfifppkikdvlmislspivtcvvvdvseddpdvqiswfvnnvevhvtaqtqthredynstlrsvsalpiqhqdwmkgkfcckvnnkdlpapiert  
iskpkgsvrapqvvyvlpppeeemtkkqvtlcmvtdfmpediyvewtngktelnykntepvlvdsdgsyfmysklrvekknwvernsyscvvhvglhnhhtksfsrtpg(471)

Predicted monomeric (non glycosylated) molecular weight: 54.4 kd. The molecule is dimeric and runs at about 120 kd in SDS-PAGE under native conditions.

**Transfectant Cell Line:** CHO

**INFORMATION:** CD274 (B7-H1, PD-L1, Programmed Death Ligand) is a member of the B7 family and is expressed on a variety of tissues including lymphoid cells. It plays an important role in regulation of T cell activation, and is involved in progression of cancer, arthritis and HIV infection (3). CD274 binding to its receptor CD279 (PD-1) on activated T cells can decrease proliferation. Conversely, ligation of CD279 on primed T cells can stimulate IL-10 production. High levels of CD274 present in Renal cell carcinoma are associated with poor prognosis (1). Tumor expressed CD274 can increase apoptosis of tumor specific T cells resulting in better tumor cell survival (2). Gamma Interferon and PHA can up regulate CD274 expression on T cells.

**REFERENCES:** 1)Cancer Research (April 2006) **66**:3381. 2)J Molecular Medicine (Feb 2004) **81**(5):281. 3)Int J Hematol (Nov 2003) 78(4):321.

**STORAGE CONDITIONS:** Store at 2 - 5°C. Open under aseptic conditions. Freeze/Thawing is not recommended.

**PRODUCT STABILITY:** Product should retain activity for at least 6 months after shipping date when stored as recommended. Ship Date:\_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl. Product was 0.1 µm filtered and vialled under aseptic conditions.

**PERFORMANCE:** Purity was >90% by SDS-PAGE. N-terminal sequence was as predicted: KPQAP. Recombinant protein was detectable in EIA by Biotinylated antibody ANC6H1 (cat #274-030) as well as clone M1H1.

*\*This Product is intended for Laboratory Research use only.*

**Ancell Corporation P.O. Box 87 243 Third Street North Bayport, MN 55003-0087 USA**  
**Phone: Toll free 800-374-9523 or 651-439-0835 Fax: 651-439-1940**