## PERFORMANCE DATA SHEET <sup>1920</sup> *Monoclonal* anti-mouse CD4/R-PE\*



*mAb name/Clone:* GK1.5 *Isotype:* Rat IgG2bк *Immunogen:* Mouse CTL-V4 membrane prep

CATALOG#: 704-050 QUANTITY: 120 tests WORKING DILUTION: 1:50 (or use 1.6µl of concentrated stock per 5 x 10<sup>5</sup>-cell test)

INFORMATION: Antibody GK1.5 binds to an extracellular epitope of the 55kd CD4 molecule present on a murine T cell subset (1,2). CD4 binds to MHC Class II expressed by antigen presenting cells, facilitating T Cell Receptor signaling. This clone has proven effective as a T cell depletion reagent(3). *References:* 1) Wilde DB, Fitch FW, et al. (1983) *J Immunol* 131(5):2178-2183. 2) Dialynas DP, Fitch FW, et al. (1983) *J Immunol* 131(5):2445-2451. 3)Mochimaru H, Ishida S, et al. (2008) *Invest Ophthamol Vis Sci* 49(5):2172-2177.

STORAGE CONDITIONS: Store at 2 - 5°C. Do not freeze! Protect from light.


**PRODUCT STABILITY:** Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:\_\_\_\_\_\_

**BUFFER:** 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 15% Glycerol, 0.2% BSA, 0.04% NaN<sub>3</sub> (as a preservative).

**PRODUCTION:** Protein G purified antibody from tissue culture supernatant was conjugated to R-Phycoerythrin through a sulfo-ester linkage. Unconjugated antibody was removed using size exclusion chromatography. Antibody conjugate is at **250 µg/ml** with an  $A_{565}/A_{280}$  ratio of 3.5.

**PERFORMANCE:** Reagent was tested for binding to ACK lysed murine splenocytes in FACS. Five x  $10^5$  splenocytes per tube were washed and pre incubated with 20 µl of 300 µg/ml Mouse IgG (to block non specific binding), after which they were incubated 45 minutes on ice with 80 µl of anti-Mouse CD4/R-PE at a 1:50 dilution (5 µg/ml). They were then washed three times, fixed and analyzed by FACS. A 27% sub population of the cells stained positive with a mean shift of 2.48  $log_{10}$  fluorescent units when compared to background. Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-mouse CD4 antibody (Catalog #704-020).

\*This Product is intended for Laboratory Research use only.



Ancell Corporation P.O. Box 87 243 Third Street North Bayport, MN 55003-0087 USA Phone: Toll free 800-374-9523 or 651-439-0835 Fax: 651-439-1940

## Binding of anti-Mouse CD4/PE to spenocytes