PERFORMANCE DATA SHEET ¹⁹²⁰ *Monoclonal* anti-mouse CD4/APC*



mAb name/Clone: GK1.5 *Isotype:* Rat IgG2bк *Immunogen:* Mouse CTL-V4 membrane prep

CATALOG#: 704-060VOLUME IN VIAL: 0.2mlQUANTITY: 120 testsVOLUME IN VIAL: 0.2mlWORKING DILUTION: 1:50 (or use 1.6μl of concentrated stock per 5 x 10⁵-cell test)

INFORMATION: Antibody GK1.5 binds to an extracellular epitope of the 55kd CD4 molecule present on a murine T cell subset (1,2). CD4 binds to MHC Class II expressed by antigen presenting cells, facilitating T Cell Receptor signaling. This clone has proven effective as a T cell depletion reagent(3). *References:* 1) Wilde DB, Fitch FW, et al. (1983) *J Immunol* 131(5):2178-2183. 2) Dialynas DP, Fitch FW, et al. (1983) *J Immunol* 131(5):2445-2451. 3)Mochimaru H, Ishida S, et al. (2008) *Invest Ophthamol Vis Sci* 49(5):2172-2177.

STORAGE CONDITIONS: Store at 2 - 5°C. Do not freeze! Protect from light.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:______


BUFFER: 50 mM Sodium Phosphate pH 7.5, 500 mM Potassium Chloride, 150mM NaCl, 15% Glycerol, 0.2% BSA, 0.04% NaN₃ (as a preservative).

PRODUCTION: Protein G purified antibody from tissue culture supernatant was conjugated to Allophycocycanin

through a sulfo-ester linkage. Unconjugated antibody was removed using size exclusion chromatography. Antibody conjugate is at **100** μ g/ml with an A₆₅₀/A₂₈₀ ratio of 2.2.

PERFORMANCE: Reagent was tested for binding to ACK lysed murine splenocytes in FACS. Five x 10^5 **splenocytes** per tube were washed and pre incubated with 20 µl of 300 µg/ml Mouse IgG (to block non specific binding), after which they were incubated 45 minutes on ice with 80 µl of anti-Mouse CD4/APC at a **1:50** dilution (2 µg/ml). They were then washed three times, fixed and analyzed by FACS. A 27% sub population of the cells stained positive with a mean shift of **2.13** log₁₀ fluorescent units when compared to background. Binding was blocked when cells were pre incubated 10 minutes with 20 µl of 0.5 mg/ml anti-mouse CD4 antibody (Catalog #704-020).

*Research use only. Not for use in Diagnostic Procedures



Ancell Corporation P.O. Box 87 Bayport, MN 55003-0087 USA Phone: Toll free 800-374-9523 or 651-439-0835 Fax: 651-439-1940

Binding of anti-Mouse CD4/APC to spenocytes